



Year: 2018

Age-predicted maximal heart rate in recreational marathon runners: a cross-sectional study on Fox's and Tanaka's equations

Nikolaïdis, Pantelis Theodoros ; Rosemann, Thomas ; Knechtle, Beat

Abstract: Age-based prediction equations of maximal heart rate (HR), such as the popular formulas Fox's 220-age, or Tanaka's $208 - 0.7 \times \text{age}$, have been widely used in various populations. Surprisingly, so far these equations have not been validated in marathon runners, despite the importance of the role of HR for training purposes in endurance running. The aim of the present study was to examine the validity of Fox and Tanaka equations in a large sample of women and men recreational marathon runners. Participants ($n = 180$, age 43.2 ± 8.5 years, VO 46.8 mL/min/kg , finishers in at least one marathon during the last year) performed a graded exercise test on a treadmill, where HR was measured. Measured HR correlated largely with age in the total sample ($r = -0.50$, $p < 0.001$), women ($r = -0.60$, $p < 0.001$) and men ($r = -0.53$, $p < 0.001$). In women, a large main effect of method on HR ($F = 0.001$, $\eta^2 = 0.294$) was shown with measured HR lower than Fox-HR (-4.8 bpm ; -8.4 , -1.3) and Tanaka-HR (-4.9 bpm ; -8.1 , -1.8). In men, a moderate effect of assessment method on HR was found ($F = 0.001$, $\eta^2 = 0.066$) with measured HR higher than Fox-HR ($+2.8$; 1.0 , 4.6), Tanaka-HR higher than Fox-HR ($+1.2$; 0.7 , 1.7). Based on these findings, it was concluded that Fox and Tanaka's formulas overestimated HR by 5 bpm in women, whereas Fox underestimated HR in men by 3 bpm. Thus, we recommend the further use of Tanaka's formula in men marathon runners. In addition, exercise physiologists and sport scientists should consider the observed differences among various assessment methods when performing exercise testing or prescribing training program relying on HR.

DOI: <https://doi.org/10.3389/fphys.2018.00226>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-152633>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Nikolaïdis, Pantelis Theodoros; Rosemann, Thomas; Knechtle, Beat (2018). Age-predicted maximal heart rate in recreational marathon runners: a cross-sectional study on Fox's and Tanaka's equations. *Frontiers in Physiology*, 9:226.

DOI: <https://doi.org/10.3389/fphys.2018.00226>



Age-Predicted Maximal Heart Rate in Recreational Marathon Runners: A Cross-Sectional Study on Fox's and Tanaka's Equations

Pantelis T. Nikolaidis¹, Thomas Rosemann² and Beat Knechtle^{2,3*}

¹ Exercise Physiology Laboratory, Nikaia, Greece, ² Institute of Primary Care, University of Zurich, Zurich, Switzerland,

³ Medbase St. Gallen Am Vadianplatz, St. Gallen, Switzerland

OPEN ACCESS

Edited by:

Luca Paolo Ardigo,
University of Verona, Italy

Reviewed by:

Brian Hanley,
Leeds Beckett University,
United Kingdom
Gustavo Zaccaria Schaun,
Universidade Federal de Pelotas,
Brazil

*Correspondence:

Beat Knechtle
beat.knechtle@hispeed.ch

Specialty section:

This article was submitted to
Exercise Physiology,
a section of the journal
Frontiers in Physiology

Received: 16 December 2017

Accepted: 28 February 2018

Published: 15 March 2018

Citation:

Nikolaidis PT, Rosemann T and
Knechtle B (2018) Age-Predicted
Maximal Heart Rate in Recreational
Marathon Runners: A Cross-Sectional
Study on Fox's and Tanaka's
Equations. *Front. Physiol.* 9:226.
doi: 10.3389/fphys.2018.00226

Age-based prediction equations of maximal heart rate (HR_{max}), such as the popular formulas Fox's 220-age, or Tanaka's $208-0.7 \times \text{age}$, have been widely used in various populations. Surprisingly, so far these equations have not been validated in marathon runners, despite the importance of the role of HR_{max} for training purposes in endurance running. The aim of the present study was to examine the validity of Fox and Tanaka equations in a large sample of women and men recreational marathon runners. Participants ($n = 180$, age 43.2 ± 8.5 years, VO_{2max} 46.8 mL/min/kg, finishers in at least one marathon during the last year) performed a graded exercise test on a treadmill, where HR_{max} was measured. Measured HR_{max} correlated largely with age in the total sample ($r = -0.50$, $p < 0.001$), women ($r = -0.60$, $p < 0.001$) and men ($r = -0.53$, $p < 0.001$). In women, a large main effect of method on HR_{max} ($p = 0.001$, $\eta^2 = 0.294$) was shown with measured HR_{max} lower than Fox- HR_{max} (-4.8 bpm; -8.4 , -1.3) and Tanaka- HR_{max} (-4.9 bpm; -8.1 , -1.8). In men, a moderate effect of assessment method on HR_{max} was found ($p = 0.001$, $\eta^2 = 0.066$) with measured HR_{max} higher than Fox- HR_{max} ($+2.8$; 1.0 , 4.6), Tanaka- HR_{max} higher than Fox- HR_{max} ($+1.2$; 0.7 , 1.7). Based on these findings, it was concluded that Fox and Tanaka's formulas overestimated HR_{max} by ~ 5 bpm in women, whereas Fox underestimated HR_{max} in men by ~ 3 bpm. Thus, we recommend the further use of Tanaka's formula in men marathon runners. In addition, exercise physiologists and sport scientists should consider the observed differences among various assessment methods when performing exercise testing or prescribing training program relying on HR.

Keywords: age groups, cardiac rate, endurance runners, graded exercise test, sex

INTRODUCTION

Training intensity and volume are predictors of performance in marathon runners (Schmid et al., 2012). A daily task in the context of training is to run at an optimal intensity in order to elicit the desired physiological adaptations, such as increased speed at anaerobic threshold and maximal oxygen uptake (Lepers and Stapley, 2016). If the intensity is inadequate, the stimulus for these adaptations is missing. On the other hand, if the intensity exceeds the optimal level, the risk of overtraining increases (O'Connor, 2007). Thus, it is important to assess exercise intensity

accurately, which relies on objective measures such as heart rate (HR), oxygen uptake and lactate, and subjective methods such as rate of perceived exertion (Foster et al., 2017). When HR is used as a measure of intensity, usually it is expressed as a function of maximal HR (HR_{max}) (Vesterinen et al., 2017).

HR_{max} can be measured using a graded exercise test (GXT) either in a laboratory or in the field (Cleary et al., 2011; Nikolaidis, 2015). However, occasionally it is not desirable to perform a GXT (e.g., to avoid the fatigue induced by maximal exercise testing close to a race or the associated financial cost). In such case, an alternative is to predict HR_{max} from an age-based equation, considering the inversely proportional relationship between age and HR_{max} . The most widely used formulas are those of Fox, Naughton, and Haskell ($HR_{max} = 220 - \text{age}$) (Fox et al., 1971) and of Tanaka, Monahan, and Seals ($HR_{max} = 208 - 0.7 \times \text{age}$) (Tanaka et al., 2001). These equations have been examined extensively in specific categories of adult population such as healthy (Nes et al., 2012), sedentary (Sarzynski et al., 2013), overweight (Franckowiak et al., 2011) and athletes (Faff et al., 2007).

Whereas the abovementioned studies have addressed many issues with regards to the validity of these popular equations of HR_{max} , there are some aspects that need further research. For instance, endurance athletes (e.g., marathon runners) and especially master athletes are under-represented in this body of research. A comparison of athletes and non-athletes had revealed lower measured- HR_{max} in the former group (Lester et al., 1968). In a recent study, it was shown that athletes of speed/power sports had similar measured- HR_{max} with endurance athletes and both had lower values than those who were untrained (Kusy and Zielinski, 2012). The decrease in HR_{max} induced by endurance training might be explained by accompanying plasma volume expansion, enhanced baroreflex function, alteration of the electrophysiology of the sinoatrial node and decreased beta-adrenergic receptor number and density (Zavorsky, 2000). Since their measured HR_{max} differs, it is reasonable to assume that the same equation of HR_{max} cannot fit in both athletes and non-athletes. This difference between athletes and non-athletes highlights the need to further examine the popular prediction equations in more samples of athletes. Considering the increasing number of those participating in marathon races (Jokl et al., 2004), the knowledge of the validity of the popular age-based equations has practical application for a large number of recreational marathon runners. Moreover, the age-based prediction of HR_{max} is a major interest for exercise physiologists when administering a GXT, where the achievement of a particular percentage of predicted HR_{max} might be necessary in order to consider the end values as maximal (Schaun, 2017). Therefore, the aim of the present study was to examine the validity of Fox- HR_{max} and Tanaka- HR_{max} in a large sample of recreational marathon runners. The research hypothesis was that these equations, which had been developed in non-athletes, would overestimate HR_{max} in recreational marathon runners due to their expected lower HR_{max} compared to non-athletes (Lester et al., 1968; Zavorsky, 2000; Kusy and Zielinski, 2012).

MATERIALS AND METHODS

Study Design and Participants

One hundred eighty-five recreational marathon runners mostly from the area of Athens volunteered to participate in this study, which had been advertised through popular websites for endurance runners. During September and October 2017, the participants visited the laboratory where they performed a GXT on a treadmill. This study was carried out in accordance with the recommendations of the Institutional Review Board of Exercise Physiology Laboratory Nikaia with written informed consent from all participants. All participants gave written informed consent in accordance with the Declaration of Helsinki. The protocol was approved by the Institutional Review Board of Exercise Physiology Laboratory Nikaia. One participant withdrew from the study during the GXT, whereas four participants did not achieve the criteria of VO_{2max} achievement, and consequently their data excluded from further analysis. Therefore, we included 180 participants from the initial sample. With regards to their sport experience, the median number of marathon completed in the past was 3 and the interquartile range was 2–6. Personal record was $4:09 \pm 0:45$ h:min.

Protocols and Equipment

Anthropometry

Height, body mass, and skinfolds were measured with participants in minimal clothing and barefoot. An electronic weighing scale (HD-351; Tanita, Arlington Heights, IL, USA) was employed for measurement of body mass (to the nearest 0.1 kg), a portable stadiometer (SECA Leicester, UK) for height (0.001 m), and a caliper (Harpender, West Sussex, UK) for skinfolds (0.2 mm). Body mass index was calculated as the quotient of body mass (kg) to height squared (m^2), and body fat (BF) was estimated from skinfolds (Parizkova, 1978).

Graded Exercise Test

A modified version of Conconi test was used to assess VO_{2max} (Conconi et al., 1982). Briefly, after a 20-min warm-up including jogging and stretching exercises, participants performed a GXT on a treadmill using a +1% inclination. The initial speed was set at 8 km/h and was increased every minute by 1 km/h till exhaustion (Christmas et al., 2017). During the late stages of the test, participants were cheered vigorously so that they made maximal effort. Measured HR_{max} was defined as the highest value attained during the test. HR was recorded continuously during the test by Team2 Pro (Polar Electro Oy, Kempele, Finland). Minute ventilation and VO_2 were recorded by a gas analyzer (Fitmate Pro, Cosmed, Rome, Italy). Anaerobic threshold was identified from ventilatory threshold, i.e., the relationship between minute ventilation and oxygen uptake. Plateau of VO_2 (primary criterion), blood lactate, age-predicted HR_{max} and RPE (secondary criteria) were used as criteria of VO_{2max} (Howley et al., 1995). Desired RPE was ≥ 8 in the 0–10 of Borg scale (Borg, 1988). Blood samples were taken 5 min after termination of test, and lactate concentration was analyzed (Accutrend, Roche, Germany). Lactate concentration was employed as a criterion of VO_{2max} achievement (accepted

values > 9 mmol/L) (Todd et al., 2017). Predicted maximal heart rate was calculated using Tanaka's formula (Tanaka et al., 2001)—as Fox's formula might overestimate HR_{max} (Nikolaidis, 2015)—and was employed as a criterion of VO₂max achievement (accepted values measured HR_{max} \geq 95% of Tanaka-HR_{max}).

Statistical Analyses

Statistical analyses were performed using IBM SPSS v.20.0 (SPSS, Chicago, IL, USA). Normality was examined using Kolmogorov-Smirnov test and visual inspection of normal Q-Q plots. Data were expressed as mean and standard deviation (SD). An independent *t*-test examined the sex differences in anthropometric and physiological characteristics. One-way repeated measures analysis of variance (ANOVA) and a subsequent Bonferroni *post-hoc* test (if there were differences among groups) were used to examine the differences between measured and predicted HR_{max}. 95% confidence intervals (CI) of the mean differences were calculated. To interpret ES for statistical differences in the ANOVA, we used eta square classified as small ($0.010 < \eta^2 \leq 0.059$), medium ($0.059 < \eta^2 \leq 0.138$), and large ($\eta^2 > 0.138$) (Cohen, 1988). Bland-Altman analysis was used to examine the accuracy and variability of prediction equations (Bland and Altman, 1986). Associations between measured HR_{max} and age were determined using Pearson's product moment correlation coefficient (*r*). Magnitude of correlation coefficients was considered as trivial if $r \leq 0.10$, small if $0.10 \leq r < 0.30$, moderate if $0.30 \leq r < 0.50$, large if $0.50 \leq r < 0.70$, very large if $0.70 \leq r < 0.90$, nearly perfect if $r \geq 0.90$, and perfect if $r = 1.00$ (Batterham and Hopkins, 2006). In addition, we used linear regression to model the prediction of HR_{max} from age in the total sample and in each sex. The linear regression was qualified for this analysis instead of non-linear regression as minimal differences among linear, quadratic, and polynomial equations have been shown (Ozemek et al., 2017). The level of significance was set at $\alpha = 0.05$.

RESULTS

The descriptive characteristics of participants are shown in Table 1. Measured HR_{max} correlated largely with age in the total sample ($r = 0.50$, $p < 0.001$), women (0.60 , $p < 0.001$) and men (0.53 , $p < 0.001$) (Figure 1). In the overall sample, no main effect of measured or predicted method on HR_{max} was observed ($p = 0.093$, $\eta^2 = 0.015$); however, *post-hoc* comparison revealed trivially larger score in Tanaka-HR_{max} than in Fox-HR_{max} (mean difference $+0.6$ bpm, $d = 0.12$), whereas no difference was found between measured HR_{max} with Fox-HR_{max} (-1.0 bpm, $d = -0.15$) and Tanaka-HR_{max} (-1.7 bpm, $d = -0.06$). A moderate sex \times assessment method interaction on HR_{max} was shown ($p < 0.001$, $\eta^2 = 0.087$) with larger differences among methods in women than in men. In women, a large main effect of method on HR_{max} ($p = 0.001$, $\eta^2 = 0.294$) was shown with measured HR_{max} lower than Fox-HR_{max} (-4.8 bpm, $d = -0.56$) and Tanaka-HR_{max} (-4.9 bpm, $d = -0.66$), and no difference between Fox-HR_{max} and Tanaka-HR_{max} (-0.1 bpm, $d = -0.01$). In men, a moderate effect of assessment

TABLE 1 | Descriptive characteristics of participants.

Parameter	Total (<i>n</i> = 180)	Women (<i>n</i> = 32)	Men (<i>n</i> = 148)
Age (years)	43.2 \pm 8.5	40.3 \pm 8.8	43.9 \pm 8.3*
Height (cm)	173.7 \pm 8.0	162.5 \pm 6.6	176.1 \pm 5.9†
Body mass (kg)	73.6 \pm 11.4	57.9 \pm 7.4	77.0 \pm 9.0†
BMI (kg.m ⁻²)	24.3 \pm 2.7	21.9 \pm 2.1	24.8 \pm 2.5†
BF (%)	18.0 \pm 4.3	19.5 \pm 4.7	17.7 \pm 4.1*
VO ₂ max (mL/min/kg)	46.8 \pm 8.9	37.5 \pm 6.7	48.8 \pm 8.1†
Lactate (mmol/L)	10.8 \pm 2.8	9.3 \pm 2.6	11.2 \pm 2.8†
RPE (a.u.)	8.8 \pm 0.9	8.4 \pm 1.0	8.8 \pm 0.8†
HR _{max} (bpm)	178.2 \pm 10.2	174.8 \pm 8.8	178.9 \pm 10.4*
Fox-HR _{max} (bpm)	176.8 \pm 8.5	179.7 \pm 8.8	176.1 \pm 8.3*
Tanaka-HR _{max} (bpm)	177.7 \pm 5.9	179.8 \pm 6.1	177.3 \pm 5.8*

BMI, body mass index; BF, body fat; VO₂max, maximal oxygen uptake; RPE, rate of perceived exertion; a.u., arbitrary units; HR_{max}, maximal heart rate; symbols *, †, and ‡ denote difference from women at $p < 0.05$, $p < 0.01$, and $p < 0.001$, respectively.

method on HR_{max} was found ($p = 0.001$, $\eta^2 = 0.066$) with measured HR_{max} higher than Fox-HR_{max} ($+2.8$ bpm, $d = 0.30$), Tanaka-HR_{max} higher than Fox-HR_{max} ($+1.2$ bpm, $d = 0.17$) and no difference between measured HR_{max} and Tanaka-HR_{max} (1.6 bpm, $d = 0.19$). The Bland-Altman plots are presented in Figure 2 (Fox-HR_{max}) and Figure 3 (Tanaka-HR_{max}). A visual inspection of these plots highlighted a trend that Tanaka-HR_{max} underestimated the low scores and overestimated the high scores in men.

DISCUSSION

The present study addressed the question whether the widely used age-based prediction equations of HR_{max}, Fox's $220 - \text{age}$ or Tanaka's $208 - 0.7 \times \text{age}$, are valid in recreation marathon runners since no study previously examined this topic. We hypothesized that these equations would overestimate HR_{max} in our sample, due to their expected lower HR_{max} compared to non-athletes (Zavorsky, 2000). The main findings were that (a) Fox-HR_{max} and Tanaka-HR_{max} overestimated HR_{max} by ~ 5 bpm in women, (b) Fox-HR_{max} underestimated HR_{max} by ~ 3 bpm in men, (c) Tanaka-HR_{max} was similar to measured-HR_{max} in men, and (d) the main effect of assessment methods on HR_{max} was larger in women than in men.

The overestimation of HR_{max} in women by age-based predicted equations was in agreement with previous findings (Esco et al., 2015). For instance, Fox and Tanaka formulas provided significantly higher estimates by 7–13 bpm compared with observed HR max in women collegiate athletes (Esco et al., 2015). Considering that the choice of assessment method had larger magnitude on women than in men, the overestimation of HR_{max} in women marathon runners is an issue that future research should address and develop sport-specific prediction equation.

The agreement between measured and Tanaka observed in men marathon runners was in line with previous research in




FIGURE 1 | Relationship between measured maximal heart rate and age.




FIGURE 2 | Bland-Altman plots of the measured maximal rate compared to the Fox's formula.




FIGURE 3 | Bland-Altman plots of the measured maximal rate compared to Tanaka's formula.

young physically active men (Barboza et al., 2016), but not with a study on sedentary adults that showed that Fox and Tanaka- HR_{max} overestimated HR_{max} in sedentary adults by 2–4 bpm (Camarda et al., 2008). Camarda et al. (2008) found that Tanaka- HR_{max} overestimated HR_{max} only by 1 bpm in men. Tanaka- HR_{max} provided closer values to HR_{max} than Fox- HR_{max} in overweight adults (Franckowiak et al., 2011) and young physically active (Barboza et al., 2016). On the other hand, Fox- HR_{max} underestimated HR_{max} in older adults (Whaley et al., 1992). In male adults, Tanaka- HR_{max} underestimated HR_{max} by 5 bpm, while there was not any difference between Fox- HR_{max} and measured- HR_{max} (Nikolaidis, 2015). Differences between the findings of the present study and those of previous research should be attributed to the chronic physiological adaptations of recreational marathon runners to endurance training. Zavorsky (2000) highlighted that endurance training results in decrease of HR_{max} due to extrinsic/autonomic (e.g., plasma volume expansion) and intrinsic/non-autonomic factors (e.g., alteration of the electrophysiology of the sinoatrial node).

The moderate sex \times assessment method interaction on HR_{max} indicated that sex should be considered in predicting HR_{max} . Women marathon runners were younger by 3.6 years and had a 4.1 bpm lower measured- HR_{max} than men indicating a relatively lower HR_{max} if sexes were age matched. This observation was in agreement with a previous study showing difference in HR_{max} among sexes (Hakki et al., 1983).

The measured HR_{max} is in agreement with previous findings on age-matched humans (Arena et al., 2016); nonetheless the variation in our sample was smaller which should be attributed to the homogeneity of the sample. On the other hand, the slopes of the linear regressions suggested that HR_{max} decreases faster in men than in women, which was in disagreement with a previous study on healthy adults showing the opposite trend (Shargal et al., 2015). An explanation of this discrepancy might be the different samples' characteristic (age and sport).

A limitation of the present study was that it focused on the prediction of HR_{max} only from age excluding other parameters that might improve the accuracy of the prediction. For instance,

Barboza et al. (2017) recommended an equation including age and HR at 150 W elicited during a GXT on a cycle ergometer in healthy young adult men. In another study, mode of exercise, fitness level, continent, and age were predictors of HR_{max} (Londeree and Moeschberger, 1982). Moreover, caution is needed to generalize the values obtained in the GXT in the laboratory to other settings, e.g., field testing, training and competition, as the latter might induce higher values (Coutinho et al., 2017). Nevertheless, strength of the present study was its novelty as it was the first to be conducted on recreational marathon runners. Considering the increasing number of those participating in marathon races, our findings are of great practical value for purposes of testing and training. Despite the different settings in laboratory and field, comparative studies observed no (Krautgasser et al., 2011; Alemдарoglu et al., 2012) or practically negligible difference (Meyer et al., 2003) in HR_{max} between these two conditions. Therefore, the findings of the present study could be applied in both laboratory and field settings, e.g., outdoors running training sessions. In addition, exercise physiologists

performing exercise testing should benefit from such knowledge in order to evaluate correctly HR as criterion of achievement of VO_{2max}.

CONCLUSIONS

Based on the present findings we recommend the further use of Tanaka's formula in men recreational marathon runners with similar training characteristics as those of the participants in the present study. In addition, exercise physiologists and sport scientists should consider the observed differences among various assessment methods when performing exercise testing or prescribing training program relying on HR.

AUTHOR CONTRIBUTIONS

PN performed the laboratory analyses, the statistical analyses and drafted the manuscript; TR and BK helped in drafting the manuscript.

REFERENCES

- Alemдарoglu, U., Dündar, U., Köklü, Y., and Asci, A. (2012). Evaluation of aerobic capacity in soccer players: comparison of field and laboratory tests. *Biol. Sport* 29, 157–161. doi: 10.5604/20831862.990468
- Arena, R., Myers, J., and Kaminsky, L. A. (2016). Revisiting age-predicted maximal heart rate: can it be used as a valid measure of effort? *Am. Heart J.* 173, 49–56. doi: 10.1016/j.ahj.2015.12.006
- Barboza, J., Nogueira, F. S., Paulucio, D., and Pompeu, F. A. M. S. (2016). The validity of maximum heart rate predictions based on age for young men. *Gazz. Med. Ital.* 175, 154–157.
- Barboza, J. A., Nogueira, F. S., and Pompeu, F. A. M. S. (2017). A new accurate model to predict maximal heart rate. *J. Exerc. Physiol. Online* 20, 23–28.
- Batterham, A. M., and Hopkins, W. G. (2006). Making meaningful inferences about magnitudes. *Int. J. Sports Physiol. Perform.* 1, 50–57. doi: 10.1123/ijsp.1.1.50
- Bland, J. M., and Altman, D. G. (1986). Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1, 307–310. doi: 10.1016/S0140-6736(86)90837-8
- Borg, G. (1988). *Borg's Perceived Exertion and Pain Scales*. Champaign, IL: Human Kinetics.
- Camarda, S. R., Tebexreni, A. S., Páfaró, C. N., Sasai, F. B., Tambeiro, V. L., Juliano, Y., et al. (2008). Comparison of maximal heart rate using the prediction equations proposed by Karvonen and Tanaka. *Arq. Bras. Cardiol.* 91, 311–314. doi: 10.1590/S0066-782X2008001700005
- Christmas, B. C. R., Taylor, L., Siegler, J. C., and Midgley, A. W. (2017). A reduction in maximal incremental exercise test duration 48 h post downhill run is associated with muscle damage derived exercise induced pain. *Front. Physiol.* 8:135. doi: 10.3389/fphys.2017.00135
- Cleary, M. A., Hetzler, R. K., Wages, J. J., Lentz, M. A., Stickley, C. D., and Kimura, I. F. (2011). Comparisons of age-predicted maximum heart rate equations in college-aged subjects. *J. Strength Cond. Res.* 25, 2591–2597. doi: 10.1519/JSC.0b013e3182001832
- Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences*, 2nd Edn. Hillsdale, NJ: Lawrence Erlbaum Associates.
- Conconi, F., Ferrari, M., Ziglio, P. G., Droghetti, P., and Codeca, L. (1982). Determination of the anaerobic threshold by a noninvasive field test in runners. *J. Appl. Physiol. Respir. Environ. Exerc. Physiol.* 52, 869–873. doi: 10.1152/jappl.1982.52.4.869
- Coutinho, C., Watson, A., Brickson, S., and Sanfilippo, J. (2017). Maximal heart rate differs between laboratory and field conditions among female athletes. *J. Hum. Sport Exerc.* 12, 386–395. doi: 10.14198/jhse.2017.122.15
- Esco, M. R., Chamberlain, N., Flatt, A. A., Snarr, R. L., Bishop, P. A., and Williford, H. N. (2015). Cross-validation of age-predicted maximal heart rate equations among female collegiate athletes. *J. Strength Cond. Res.* 29, 3053–3059. doi: 10.1519/JSC.0000000000000978
- Faff, J., Sitkowski, D., Ladyga, M., Klusiewicz, A., Borkowski, L., and Starczewska-Czapowska, J. (2007). Maximal heart rate in athletes. *Biol. Sport* 24, 129–142.
- Foster, C., Rodriguez-Marroyo, J. A., and de Koning, J. J. (2017). Monitoring training loads: the past, the present, and the future. *Int. J. Sports Physiol. Perform.* 12, S22–S28. doi: 10.1123/IJSP.2016-0388
- Fox, S. M. III., Naughton, J. P., and Haskell, W. L. (1971). Physical activity and the prevention of coronary heart disease. *Ann. Clin. Res.* 3, 404–432.
- Franckowiak, S. C., Dobrosielski, D. A., Reilley, S. M., Walston, J. D., and Andersen, R. E. (2011). Maximal heart rate prediction in adults that are overweight or obese. *J. Strength Cond. Res.* 25, 1407–1412. doi: 10.1519/JSC.0b013e3181d682d2
- Hakki, A. H., Hare, T. W., and Iskandrian, A. S. (1983). Prediction of maximal heart rates in men and women. *Cardiovasc. Rev. Rep.* 4, 997–999.
- Howley, E. T., Bassett, D. R. Jr., and Welch, H. G. (1995). Criteria for maximal oxygen uptake: review and commentary. *Med. Sci. Sports Exerc.* 27, 1292–1301. doi: 10.1249/00005768-199509000-00009
- Jokl, P., Sethi, P. M., and Cooper, A. J. (2004). Master's performance in the New York City Marathon 1983–1999. *Br. J. Sports Med.* 38, 408–412. doi: 10.1136/bjsm.2002.003566
- Krautgasser, S., Scheiber, P., von Duvillard, S. P., and Müller, E. (2011). Physiological responses of elderly recreational alpine skiers of different fitness and skiing abilities. *J. Sports Sci. Med.* 10, 748–753.
- Kusy, K., and Zielinski, J. (2012). Aerobic capacity in speed-power athletes aged 20–90 years vs endurance runners and untrained participants. *Scand. J. Med. Sci. Sports* 24, 68–79. doi: 10.1111/j.1600-0838.2012.01496.x
- Lepers, R., and Stapley, P. J. (2016). Master athletes are extending the limits of human endurance. *Front. Physiol.* 7:613. doi: 10.3389/fphys.2016.00613
- Lester, M., Sheffield, L. T., Trammell, P., and Reeves, T. J. (1968). The effect of age and athletic training on the maximal heart rate during muscular exercise. *Am. Heart J.* 76, 370–376. doi: 10.1016/0002-8703(68)90233-0
- Londeree, B. R., and Moeschberger, M. L. (1982). Effect of age and other factors on maximal heart rate. *Res. Q. Exerc. Sport* 53, 297–304. doi: 10.1080/02701367.1982.10605252
- Meyer, T., Welter, J.-P., Scharhag, J., and Kindermann, W. (2003). Maximal oxygen uptake during field running does not exceed that measured during treadmill exercise. *Eur. J. Appl. Physiol.* 88, 387–389. doi: 10.1007/s00421-002-0718-x

- Nes, B. M., Janszky, I., Wisløff, U., Støylen, A., and Karlsen, T. (2012). Age-predicted maximal heart rate in healthy subjects: the HUNT fitness study. *Scand. J. Med. Sci. Sports* 23, 697–704. doi: 10.1111/j.1600-0838.2012.01445.x
- Nikolaidis, P. T. (2015). Maximal heart rate in soccer players: measured versus age-predicted. *Biomed. J.* 38, 84–89. doi: 10.4103/2319-4170.131397
- O'Connor, P. J. (2007). Monitoring and titrating symptoms: a science-based approach to using your brain to optimise marathon running performance. *Sports Med.* 37, 408–411. doi: 10.2165/00007256-200737040-00035
- Ozemek, C., Whaley, M. H., Finch, W. H., and Kaminsky, L. A. (2017). Maximal heart rate declines linearly with age independent of cardiorespiratory fitness levels. *Eur. J. Sport Sci.* 17, 563–570. doi: 10.1080/17461391.2016.1275042
- Parizkova, J. (1978). "Lean body mass and depot fat during autogenesis in humans," in *Nutrition, Physical Fitness and Health: International Series on Sport Sciences*, eds J. Parizkova and V. Rogozkin (Baltimore: University Park Press), 20.
- Sarzynski, M. A., Rankinen, T., Earnest, C. P., Leon, A. S., Rao, D. C., Skinner, J. S., et al. (2013). Measured maximal heart rates compared to commonly used age-based prediction equations in the heritage family study. *Am. J. Hum. Biol.* 25, 695–701. doi: 10.1002/ajhb.22431
- Schaun, G. Z. (2017). The maximal oxygen uptake verification phase: a light at the end of the tunnel? *Sports Med. Open* 3:44. doi: 10.1186/s40798-017-0112-1
- Schmid, W., Knechtle, B., Knechtle, P., Barandun, U., Rüst, C. A., Rosemann, T., et al. (2012). Predictor variables for marathon race time in recreational female runners. *Asian J. Sports Med.* 3, 90–98. doi: 10.5812/asjms.34704
- Shargal, E., Kislev-Cohen, R., Zigel, L., Epstein, S., Pilz-Burstein, R., and Tenenbaum, G. (2015). Age-related maximal heart rate: Examination and reinement of prediction equations. *J. Sports Med. Phys. Fitness* 55, 1207–1218.
- Tanaka, H., Monahan, K. D., and Seals, D. R. (2001). Age-predicted maximal heart rate revisited. *J. Am. Coll. Cardiol.* 37, 153–156. doi: 10.1016/S0735-1097(00)01054-8
- Todd, J. J., McSorley, E. M., Pourshahidi, L. K., Madigan, S. M., Laird, E., Healy, M., et al. (2017). Vitamin D3 supplementation using an oral spray solution resolves deficiency but has no effect on VO2 max in Gaelic footballers: results from a randomised, double-blind, placebo-controlled trial. *Eur. J. Nutr.* 56, 1577–1587. doi: 10.1007/s00394-016-1202-4
- Vesterinen, V., Nummela, A., Laine, T., Hynynen, E., Mikkola, J., and Häkkinen, K. (2017). A submaximal running test with postexercise cardiac autonomic and neuromuscular function in monitoring endurance training adaptation. *J. Strength Cond Res.* 31, 233–243. doi: 10.1519/JSC.0000000000001458
- Whaley, M. H., Kaminsky, L. A., Dwyer, G. B., Getchell, L. H., and Norton, J. A. (1992). Predictors of over- and underachievement of age-predicted maximal heart rate. *Med. Sci. Sports Exerc.* 24, 1173–1179. doi: 10.1249/00005768-199210000-00017
- Zavorsky, G. S. (2000). Evidence and possible mechanisms of altered maximum heart rate with endurance training and tapering. *Sports Med.* 29, 13–26. doi: 10.2165/00007256-200029010-00002

Conflict of Interest Statement: BK was employed by Medbase St. Gallen Am Vadianplatz.

The other authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Nikolaidis, Rosemann and Knechtle. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.